36th International Mathematical Olympiad

Canada, July 1995.

1. Let A, B, C, D be distinct points on a line, in that order. The circles with diameters AC and BD intersect at X and Y. O is an arbitrary point on the line XY but not on AD. CO intersects the circle with diameter AC again at M, and BO intersects the other circle again at N. Prove that AM, DN and XY are concurrent.

Soln. The result is trivial if O coincides with X or Y. Suppose it does not. From

$$ON \cdot ON = OX \cdot OY = OC \cdot OM$$
,

BCMN is cyclic. If O is on the segment XY, then

$$\angle MAD + \angle MNB + \angle BND = \angle MAD + \angle MCA + \angle AMC = 180^{\circ} >$$


If O is not on the segment XY, then

$$\angle MAD = 180^{\circ} - \angle AMC - \angle MAC = 180^{\circ} - \angle BND - \angle ONM = \angle MND.$$

In either case, ADNM is also cyclic. Let AM and DN intersect at Z. Let the line ZX intersect the first circle at Y_1 and the second at Y_2 . Then

$$ZX \cdot ZY_1 = ZA \cdot ZM = ZD \cdot ZN = ZX \cdot ZY_2.$$

Hence $Y_1 = Y_2 = Y$ and indeed Z lies on XY.



2. Let a, b and c be positive real numbers such that abc = 1. Prove that

$$\frac{1}{a^3(b+c)} + \frac{1}{b^3(a+b)} + \frac{1}{c^3(a+b)} \ge \frac{3}{2}.$$

Soln. Let S be the right hand side and $T = a(b+c) + b(a+c) + c(a+b) = 2(ab+ac+bc) \ge 6(abc)^{2/3} = 6$. Then by Cauchy's inequality, we have

$$ST \ge \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)^2 = \frac{T^2}{4}.$$

Thus $S \geq T/4 \geq 3/2$ as desired.

Second soln. Let x = 1/a. y = 1/b, z = 1/c. Then xyz = a and

$$S = \frac{x^2}{y+z} + \frac{y^2}{x+z} + \frac{z^2}{x+y}.$$

By Cauchy's inequality,

$$[(x+y) + (z+x) + (x+y)]S \ge (x+y+z)^2$$

or $S \ge \frac{(x+y+z)}{2} \ge \frac{3}{2}(xyz)^{1/3} = \frac{3}{2}$ as desired.

- **3.** Determine all integers n > 3 such that there are n points A_1, A_2, \ldots, A_n in the plane which satisfy the following two conditions simultaneously.
- (a) No three lie on the same line.
- (b) There exist real numbers p_1, p_2, \ldots, p_n such that the area of $\triangle A_i A_j A_k$ is equal to $p_i + p_j + p_k$ for $1 \le i < j < k \le n$.

Soln. We claim n=4 is the only answer. For n=4, let $A_1A_2A_3A_4$ be a unit square and let $p_1=p_2=p_3=p_4=1/6$. It remains to show no solution exists for n=5 which implies that there no solution for any $n\geq 5$.

Suppose to the contrary that there is solution with n=5. Denote the area of $\triangle A_i A_j A_k$ by $[ijk] = p_i + p_j + p_k$. We cannot have $p_i = p_j$. if for example $p_4 = p_5$, then [124] = [125] and [234] = [235], which implies that $A_4 A_5$ is parallel to both $A_1 A_2$ and $A_2 A_3$. This is impossible since A_1, A_2, A_3 are not collinear. We also observe that if $A_i A_j A_k A_l$ is convex, then $p_i + p_k = p_j + p_l$. This follows from [ijk] + [kli] = [jkl] + [lij].

Consider the convex hull of A_1, \ldots, A_5 , we have 3 cases. First, suppose that the convex hull is a pentagon $A_1A_2A_3A_4A_5$. Since $A_1A_2A_3A_4$ and $A_1A_2A_3A_5$ are convex, our observation yields $p_1 + p_3 = p_2 + p_4$ and $p_1 + p_3 = p_2 + p_5$. Hence $p_4 = p_5$, a contradiction.

Next we suppose that the convex hull is a quadrilateral $A_1A_2A_3A_4$. We may assume that A_5 is within $A_3A_4A_1$. Then $A_1A_2A_3A_5$ is convex and we have the same contradiction as before.

Finally suppose that the convex hull is a triangle $A_1A_2A_3$. Since [124] + [234] + [314] = [125] + [235] + [315] we have $p_4 = p_5$, a contradiction.

4. The positive real numbers $x_0, x_1, \ldots, x_{1995}$ satisfy $x_0 = x_{1995}$ and

$$x_{i-1} + \frac{2}{x_{i-1}} = 2x_i + \frac{1}{x_i}$$

for i = 1, 2, ..., 1995. Find the maximum value that x_0 can have.

Soln. The given condition is equivalent to

$$(2x_i^2 - x_{i-1})(x_i - \frac{1}{x_{i-1}}) = 0$$

which yield $x_i = x_{i-1}/2$ and $x_i = 1/x_{i-1}$. We shall prove by induction that $x_i = 2^{k_i} x_0^{\epsilon_i}$, where $|k_i| \leq i$ and $\epsilon_i = (-1)^{k_i+i}$. This is true for i = 0, with $k_0 = 0$, $\epsilon_0 = 1$. Assume that it is true for i - 1. Then for $x_i = \frac{1}{2}x_{i-1}$, we have $k_i = k_{i-1} - 1$ and $\epsilon_i = \epsilon_{i-1}$. For $x_i = 1/x_{i-1}$, we have $k_i = -k_{i-1}$ and $\epsilon_i = -\epsilon_{i-1}$. In each case, it is immediate that $|k_i| \leq i$ and $\epsilon_i = (-1)^{k_i+i}$. Thus $x_0 = x_{1995} = 2^k x_0^{\epsilon}$ where $k = k_{1995}$ and $\epsilon = \epsilon_{1995}$. Thus $|k| \leq 1995$ and $\epsilon = (-1)^{1995+k}$. If k is odd then $\epsilon = 1$ and this is impossible. Thus k is even and $\epsilon = -1$ and $x_0^2 = 2^k$. Since $|k| \leq 1995$, we have $k \leq 1994$. Hence $x_0 \leq 2^{997}$. The value $x_0 = 2^{997}$ can be attained by choosing $x_i = \frac{1}{2}x_{i-1}$ for $i = 1, 2, \ldots, 1994$ and $x_{1995} = 1/x_{1994}$.

5. Let ABCDEF be a convex hexagon with AB = BC = CD, DE = EF = FA and $\angle BCD = \angle EFA = \pi/3$. Let G and H be two points interior to the hexagon such that angles AGB and DHE are both $2\pi/3$. Prove that $AG + GB + GH + DH + HE \ge CF$.

Soln. Note that BCD and EFA are equilateral triangles. Hence BE is an axis of symmetry of ABDE. Reflect BCD and EFA about BE to BC'A and EF'D respectively. Since $\angle BGA = 180^{\circ} - \angle AC'B$, G lies on the circumcircle of ABC'. By Ptolemy's Theorem, AG + GB = C'G. Similarly, DH + HE = HF'. It follows that

$$CF = C'F' \le C'G + GH + HF' = AG + GB + GH + DH + HE,$$

with equality if and only if G and H both lie on C'F'.

6. Let p be an odd prime. Find the number of p-element subsets A of $\{1, 2, \ldots, 2p\}$ such that the sum if all elements of A is divisible by p.

Soln. For any p-element subset A of $\{1, 2, ..., 2p\}$, denote by s(A) the sum of the elements of A. Of the $\binom{2p}{p}$ such subsets, $B = \{1, 2, ..., p\}$ and $C = \{p + 1, p + 2, ..., 2p\}$ satisfy $s(B) = s(C) \equiv 0 \pmod{p}$. For $A \neq B, C$, we have $A \cap B \neq \emptyset \neq A \cap C$. Partition the $\binom{2p}{p} - 2$ p-element subsets other than B and C into groups of size p as follows. Two subsets A and A' are in the same group if and only if $A' \cap C = A \cap C$ and $A' \cap B$ is a cyclic

permutation of $A \cap B$ within B. Suppose $A \cap B$ has n elements, 0 < n < p. For some m such that 0 < m < p,

$$A' \cap B = \{x + m : x \in A \cap B, x + m \le p\} \cup \{x + m - p : x \in A \cap B, x \le p < x + m\}.$$

Hence $s(A') - s(A) \equiv mn \pmod{p}$, but mn is not divisible by p. It follows that exactly one subset A in each group satisfies $s(A) \equiv 0 \pmod{p}$, and the total number of such subsets is $p^{-1}\left(\binom{2p}{p}-2\right)+2$.

Second soln. Let ω be a primitive pth root of unity. Then

$$a_0 + a_1\omega + \dots + a_{p-1}\omega^{p-1} = 0$$
 iff $a_0 = a_1 = \dots = a_{p-1}$.

Also

$$\prod_{k=1}^{2p} (x - \omega^k) = (x^p - 1)^2 = x^{2p} - 2x^p + 1.$$

If $t(\omega) = \sum a_i \omega^i$, then a_i is the number of combinations with $i_1 + \cdots + i_p \equiv i \pmod{p}$. Therefore

$$(a_0 - 2) + a_1\omega + \dots + a_{p-1}^{p-1} \equiv 0$$

and
$$a_0 - 2 = a_1 = \dots = a_p$$
. So $|S| = \frac{\binom{2p}{p} - 2}{p} + 2$.