## **CWMI 2018**

15 August 2018

- 1. Real numbers  $x_1, x_2, \ldots, x_{2018}$  satisfy  $x_i + x_j \ge (-1)^{i+j}$  for all  $1 \le i < j \le 2018$ . Find the minimum possible value of  $\sum_{i=1}^{2018} ix_i$ .
- 2. Let  $n \ge 2$  be an integer. Positive reals  $x_1, x_2, \dots, x_n$  satisfy  $x_1 x_2 \dots x_n = 1$ . Show:

$$\{x_1\} + \{x_2\} + \dots + \{x_n\} < \frac{2n-1}{2}$$

Where  $\{x\}$  denotes the fractional part of x.

- 3. Let  $M = \{1, 2, \dots, 10\}$ , and let T be a set of 2-element subsets of M. For any two different elements  $\{a, b\}, \{x, y\}$  in T, the integer (ax + by)(ay + bx) is not divisible by 11. Find the maximum size of T.
- 4. In acute angled  $\triangle ABC$ , AB > AC, points E, F lie on AC, AB respectively, satisfying BF + CE = BC. Let  $I_B, I_C$  be the excenters of  $\triangle ABC$  opposite B, Crespectively,  $EI_C, FI_B$  intersect at T, and let K be the midpoint of arc BAC. Let KT intersect the circumcircle of  $\triangle ABC$  at K, P. Show T, F, P, E concyclic.

16 August 2018

- 5. In acute triangle ABC, AB < AC, O is the circumcenter of the triangle. M is the midpoint of segment BC, (AOM) intersects the line AB again at D and intersects the segment AC at E. Prove that DM = EC.
- 6. Let  $n \ge 2$  be an integer. Positive reals satisfy  $a_1 \ge a_2 \ge \cdots \ge a_n$ . Prove that  $\left(\sum_{i=1}^n \frac{a_i}{a_{i+1}}\right) n \le \frac{1}{2a_1a_n} \sum_{i=1}^n (a_i a_{i+1})^2$ , where  $a_{n+1} = a_1$ .
- 7. Let p and c be an prime and a composite, respectively. Prove that there exist two integers m, n, such that  $0 < m n < \frac{\operatorname{lcm}(n+1,n+2,\cdots,m)}{\operatorname{lcm}(n,n+1,\cdots,m-1)} = p^c$ .
- 8. Let n, k be positive integers, satisfying n is even,  $k \ge 2$  and n > 4k. There are n points on the circumference of a circle. If the endpoints of  $\frac{n}{2}$  chords in a circle that do not intersect with each other are exactly the n points, we call these chords a matching.Determine the maximum of integer m, such that for any matching, there exists k consecutive points, satisfying all the endpoints of at least m chords are in the k points.